

M.L. 2015 Project Abstract

For the Period Ending June 30, 2018

PROJECT TITLE: A Foundational dataset characterizing historic forest disturbance patterns

PROJECT MANAGER: Michael Falkowski

AFFILIATION: University of Minnesota – Department of Forest Resources and Cloquet Forestry Center

MAILING ADDRESS: 175 University Road

CITY/STATE/ZIP: Cloquet, MN 55720

PHONE: (208) 218-726-6410

E-MAIL: mfalkows@umn.edu

WEBSITE: <http://cfc.cfans.umn.edu/index.htm>

FUNDING SOURCE: Environment and Natural Resources Trust Fund

LEGAL CITATION: M.L. 2015, Chp. 76, Sec. 2, Subd. 03q

APPROPRIATION AMOUNT: \$ 200,000

AMOUNT SPENT: \$ 200,000

AMOUNT REMAINING: \$ 0.00

Overall Project Outcome and Results

Forest disturbance (arising from harvesting, fire, land conversion, etc.) has a fundamental impact on the health and resilience of multiple forest resources including water quality, wildlife habitat, and wood resources, among others. Recently the United States Geological Survey made a revolutionary decision by allowing open access to a historic archive of Landsat satellite data dating back to 1972, providing a new opportunity to assess historic forest disturbance (type, timing, and patterns). The objective of this project was to utilize the historical satellite images to characterize >40 years of Minnesota forest trends and disturbance patterns, and provide spatial mapping resources for a variety of local forest management and research applications. After the necessary processing to compile the Landsat imagery in a way that would allow the data to be comparable through time, we created models to produce annual (1973-2015) state-wide maps of canopy cover. These maps allow for the characterization of forest resources at a given point in time, as well as the monitoring of forest change and recovery trends, providing a valuable and versatile dataset for a variety of Minnesota users. For the second part of this project, we focused on the Laurentian Mixed Forest Province, which contains much of the public and forested lands of Minnesota, where we utilized additional Landsat data to map the most recent abrupt disturbance events over time. We further enhanced the disturbance map by classifying the disturbance agent (harvest, land conversion, fire, wind, flooding), as well as providing information about the year, duration, and magnitude of each event. Currently we are working with several collaborators to input our mapping products to address a variety of forest management, wildlife habitat, and water quality assessment applications.

Project Results Use and Dissemination

Our initial publication from this project, entitled “Extracting the full value of the Landsat archive: Inter-sensor harmonization for the mapping of Minnesota forest canopy cover (1973–2015)” was published in Remote Sensing and Environment in March 2018, and is already providing a valuable resource for fellow researchers through our approach for incorporating rarely integrated early Landsat MSS imagery to time series analyses for the creation of >40 years of annual forest attribute mapping. While only recently published, the paper has already received 4 citations in peer reviewed publications and boasts 423 reads on research focused social media platform. We were invited to present this work through a webinar for the USDA Forest Service’s Forest Inventory and Analyses National Research Techniques Band (recording

available at: <https://usfs.adobeconnect.com/prjhzov1f5fi/>), and continue to utilize the valuable state-wide data set presented in this publication for our disturbance mapping efforts and various forest, wildlife, and water resources applications.

We have worked with, and continue to work with, several collaborators to provide our canopy cover and disturbance mapping products for a variety of forest management, wildlife habitat, and water quality assessment applications. In addition to providing mapping resources to UMN moose biologists to assess habitat use and movement, we are also currently working with wildlife researchers from UMN-Duluth to incorporate our canopy cover and disturbance mapping products in a project assessing the impacts of harvest intensities and the quantity and spatial arrangement of retained tree canopy on avian and small mammal communities across a chrono-sequence of harvest ages. We have also provided initial harvest maps to contractors working with the MN PCA, to incorporate into a watershed planning tool for assessing forestry best management practices and impacts on water quality.

We have presented our work to a variety of research groups, local managers, and state and federal agencies throughout the project time period, and we continue to disseminate our results and mapping products to a variety of audiences to ensure that our products can provide vital additions to existing projects and management planning needs. We also continue to explore additional applications of the data and are working to compile manuscripts related to utilizing the disturbance products to assess various forest ecology and resource management questions and issues.

Date of Report: 8-17-2018

Final Report

Date of Work Plan Approval: 06-11-2015

Project Completion Date: 06-30-2018

PROJECT TITLE: A foundational dataset characterizing historic forest disturbance patterns

Project Manager: Michael Falkowski

Organization: University of Minnesota – Department of Forest Resources and Cloquet Forestry Center

Mailing Address: 175 University Road

City/State/Zip Code: Colquet, MN 55720

Telephone Number: (208) 218-726-6410

Email Address: mfalkows@umn.edu

Web Address: <http://cfc.cfans.umn.edu/index.htm>

Location: Carlton, Lake, and St. Louis Counties

Total ENRTF Project Budget:

ENRTF Appropriation: \$200,000

Amount Spent: \$200,000

Balance: \$0

Legal Citation: M.L. 2015, Chp. 76, Sec. 2, Subd. 03q

Appropriation Language:

\$200,000 the first year is from the trust fund to the Board of Regents of the University of Minnesota to quantify forest disturbance impacts over the past forty years on water quality, wildlife demographics, and wood fiber supply to identify management strategies that better respond to disturbance impacts and improve and sustain forest resources. This appropriation is available until June 30, 2018, by which time the project must be completed and final products delivered.

I. PROJECT TITLE:

A foundational dataset characterizing historic forest disturbance patterns

II. PROJECT STATEMENT:

Forest disturbance dynamics (arising from harvesting, fire, land conversion, etc.) have a fundamental control on the health and resilience of multiple forest resources including water quality, wildlife habitat, and wood resources, among others. Disturbance impacts on forest resources can persist across a landscape for decades and ultimately impact the sustainability of resources, positively or negatively. Understanding disturbance dynamics and associated impacts is readily recognized as being critically important to developing forest management responses that improve and sustain forest resources. Despite this recognition, until recently it has been nearly impossible to quantify and interpret disturbance configurations (type, timing, and pattern) that have persistent impacts on forest resources. Recently the United States Geological Survey made a revolutionary decision by allowing open access to a 40-year historic archive of Landsat satellite data, providing a new opportunity to assess historic forest disturbance dynamics. These satellite-derived disturbance observations can be used to (i) determine the fundamental drivers of past disturbance and (ii) assess the impacts of disturbances on current forest resources.

The overarching objective of this project is to leverage historical satellite images in the archive to develop a foundational dataset that characterizes trends and patterns in historic forest disturbances Across Minnesota's arrowhead region. As a secondary objective we will demonstrate the utility of foundational disturbance dataset to sustainable resource management in two key areas: wildlife habitat and wood resource management.

We will develop the foundational dataset characterizing trends and patterns in forest disturbance by processing and analyzing Landsat satellite images (1974-present) from the aforementioned archive. This step will involve: (i) acquiring cloud free imagery across each growing season from 1974-present, (ii) preprocessing the imagery to ensure intra-seasonal and inter-annual comparability of the images, and (iii) running semi-automated image classification procedures that characterize disturbance type (e.g., fire, insect outbreak, blow-down, etc.) and duration of impacts and ecosystem recovery (e.g., how long did the disturbance last and how long did it take the ecosystem to return to its pre-disturbance state). The utility of this dataset to resource management will be demonstrated in specific applied examples. For the case of wildlife habitat we will integrate disturbance data with historic demographic data characterizing moose populations across Minnesota's Arrowhead region to understand how disturbance can enhance or degrade moose habitat quality and ultimately population sustainability. For the wood resource example we will link information from the disturbance dataset with repeated forest inventory data (e.g., data from the US forest service FIA (Forest Inventory and Analysis) program) to understand the impacts of different disturbance types on wood resources across Minnesota's Arrowhead region. In this example we will specifically develop an understanding how disturbances impact the amount (e.g., biomass, volume, etc.) and quality (e.g., species composition, age, etc.) of the forest resource landscape, which ultimately impact long term resource sustainability.

The foundational dataset will be a powerful tool for identifying threshold disturbance patterns that positively or negatively impact multiple forest resources across Minnesota. This foundational dataset will allow end users to evaluate how the type, timing, duration, and configuration of disturbances influence forest resources over the last forty years, and will ultimately help in the identification of management responses that improve and sustain forest resources into the future.

III. OVERALL PROJECT STATUS UPDATES:

Project Status as of February 2016:

The project is moving forward successfully. All appropriate Landsat growing season images have been selected, requested, and downloaded for the time period of 1972-2015 and much of the pre-processing completed. Methods were developed and implemented for the initial pre-processing steps, and the next steps have been outlined and in some cases tested with small subsets of data (including the next immediate step of calibrating

the disturbance trend algorithm, LandTrendr). Pre-processing work is well underway for all scenes including the unpacking of data and initial quality assessments of images, geo-referencing of images where needed, calculation of spectral indices, calibration between Landsat sensors (MSS, TM, and OLI), and compositing of images into annual composites of growing season spectral indices used for later disturbance/recovery algorithms and classification. Initial contact has been made with potential Minnesota wildlife research collaborators (including Jerry Niemi's lab group and MN DNR Wildlife Habitat team), to solicit input on their specific needs for disturbance products and initiate collaborative project planning.

Project Status as of *September 2016*:

After substantial work in the pre-processing of Landsat images to produce corrected, calibrated, and comparable annual stacks of spectral information, we now have usable annual growing season composites for the entire Arrowhead region from 1972-2015. We have continued forward with these image stacks, completing the first step of the change detection algorithm, LandTrendr, which segments the spectral trends through time and assigns smoothed values important in identifying disturbance and recovery patches and providing information on the timing, duration, and magnitude of change. To provide initial insight into historic forest cover and change and to provide information about an important wildlife habitat component through time, we are currently creating models for canopy cover which will be mapped across the 44 year image stack. Forest Inventory and Analysis data and plot locations along with two years of NAIP imagery has been obtained for the study area in preparation for building training and validation data sets of known disturbance agents vital to the future steps of the project where we will classify and map disturbance agent and characteristics of recovery trends.

Project Status as of *February 2017*:

As initial products important to a variety of applications including wildlife habitat modeling, forest monitoring, and to aid in the next steps of identifying and interpreting forest change and recovery, we have finish the creation of forest mask and continuous canopy cover models for the Arrowhead region. This involved NAIP photo interpretation of canopy cover at >1000 FIA plot locations for consistent estimates of cover across all land cover types, and exploration into several statistical modeling approaches in order to determine the most appropriate method for minimizing canopy cover prediction error. In addition to the model for continuous canopy cover, we also developed a forest mask model which can differentiate forest from all other land cover types, a vital tool for next steps of the project, as well as for monitoring trends of forest area change over the last 4 decades. This forest mask and canopy cover models are in the final stage of creating predictive maps using the 44 years of Landsat imagery, which will result in stacks of annual forest masks and continuous canopy cover maps for the entire Arrowhead Region across all land cover types from 1972-2015. We plan on submitting the manuscript pertaining to this work for peer review in the coming months. We are also actively planning for immediate incorporation of these products in a variety of applications once complete.

Project Status as of *September 2017*:

Following completion of the annual canopy cover and forest mask maps (1973-2015) described above, we have prepared the associated manuscript and submitted to Remote Sensing of Environment. The manuscript also includes many of the preprocessing steps for our larger change detections work, including the compiling of the Landsat stacks into annual harmonized composites of the spectral indices utilizing a new R package called LandsatLinkr which we beta-tested in this project. Initial reviews of the manuscript were favorable and we are currently addressing revisions. Successful publication of this work will serve as an important introduction of methods for our subsequent change detection manuscripts as well as providing insight into Minnesota forest trends across the last four decades. In addition to our canopy cover mapping manuscript work, we have completed initial runs of LandTrendr change detection although we continue to tweak parameters to best match the disturbance patterns in the Arrowhead Regions and are working to compile the agent classification training data base. Preliminary exploration into post-processing of the change products for the mapping of true change

polygons of fast and slow disturbances and classifications of agent of change are currently underway. We have also conducted field visits to various locations around the Arrowhead region which included meetings with foresters within the Chippewa National Forest, Superior National Forest, and the UMN Cloquet Forestry Center to discuss local disturbance patterns, future applications of mapping products, and the needs of the on-the-ground forest managers.

Project Status as of *February 2018*:

Our manuscript presenting our initial methods for producing harmonized spectral indices for the Landsat Time Series, and the application of the data for the creation of annual canopy cover and forest mask maps from 1973-2015 for the state of Minnesota, has been accepted for publication within *Remote Sensing of Environment*. On the disturbance mapping front, we continue to do quality assessments of the initial change detection products to finalize the parameters utilized in the identification of disturbance patches across the Arrowhead Region. Change mapping is a multi-iterative process, where we model and map change patches and create classification of agents using training data, assess the maps, and evaluate where additional training data, predictive metrics, and/or calibration of parameters may be needed for the next iteration of the process. With initial models and maps complete, we are currently working to improve the quality of the data sets to work towards finalized disturbance products and validation assessments. We were able to present our initial findings and our overall project methods and products at multiple meetings within the twin cities and Cloquet in January 2018. These meetings included presenting an invited webinar to members of MN DNR, forest service, and UMN affiliates; participating in, and presenting at the 2018 SFEC Forestry and Wildlife Research Review meeting at the Cloquet Forestry Center; and multiple smaller meetings with various forest service, MN DNR, and UMN researchers to discuss potential collaborations and applications of our data products. All of these meetings displayed a wide breadth of interest from a variety of research and management agencies in the use of our forest attribute and change detection products for a range of forest, wildlife, water quality, and monitoring applications.

Overall Project Outcomes and Results:

Forest disturbance (arising from harvesting, fire, land conversion, etc.) has a fundamental impact on the health and resilience of multiple forest resources including water quality, wildlife habitat, and wood resources, among others. Recently the United States Geological Survey made a revolutionary decision by allowing open access to a historic archive of Landsat satellite data dating back to 1972, providing a new opportunity to assess historic forest disturbance (type, timing, and patterns). The objective of this project was to utilize the historical satellite images to characterize >40 years of Minnesota forest trends and disturbance patterns, and provide spatial mapping resources for a variety of local forest management and research applications. After the necessary processing to compile the Landsat imagery in a way that would allow the data to be comparable through time, we created models to produce annual (1973-2015) state-wide maps of canopy cover. These maps allow for the characterization of forest resources at a given point in time, as well as the monitoring of forest change and recovery trends, providing a valuable and versatile dataset for a variety of Minnesota users. For the second part of this project, we focused on the Laurentian Mixed Forest Province, which contains much of the public and forested lands of Minnesota, where we utilized additional Landsat data to map the most recent abrupt disturbance events over time. We further enhanced the disturbance map by classifying the disturbance agent (harvest, land conversion, fire, wind, flooding), as well as providing information about the year, duration, and magnitude of each event. Currently we are working with several collaborators to input our mapping products to address a variety of forest management, wildlife habitat, and water quality assessment applications.

Project Results Use and Dissemination:

Our initial publication from this project, entitled "Extracting the full value of the Landsat archive: Inter-sensor harmonization for the mapping of Minnesota forest canopy cover (1973–2015)" was published in *Remote Sensing and Environment* in March 2018, and is already providing a valuable resource for fellow researchers through our approach for incorporating rarely integrated early Landsat MSS imagery to time series analyses for

the creation of >40 years of annual forest attribute mapping. While only recently published, the paper has already received 4 citations in peer reviewed publications and boasts 423 reads on research focused social media platform. We were invited to present this work through a webinar for the USDA Forest Service’s Forest Inventory and Analyses National Research Techniques Band (recording available at: <https://usfs.adobeconnect.com/prjhzov1f5fi/>), and continue to utilize the valuable state-wide data set presented in this publication for our disturbance mapping efforts and various forest, wildlife, and water resources applications.

We have worked with, and continue to work with, several collaborators to provide our canopy cover and disturbance mapping products for a variety of forest management, wildlife habitat, and water quality assessment applications. In addition to providing mapping resources to UMN moose biologists to assess habitat use and movement, we are also currently working with wildlife researchers from UMN-Duluth to incorporate our canopy cover and disturbance mapping products in a project assessing the impacts of harvest intensities and the quantity and spatial arrangement of retained tree canopy on avian and small mammal communities across a chrono-sequence of harvest ages. We have also provided initial harvest maps to contractors working with the MN PCA, to incorporate into a watershed planning tool for assessing forestry best management practices and impacts on water quality.

We have presented our work to a variety of research groups, local managers, and state and federal agencies throughout the project time period, and we continue to disseminate our results and mapping products to a variety of audiences to ensure that our products can provide vital additions to existing projects and management planning needs. We also continue to explore additional applications of the data and are working to compile manuscripts related to utilizing the disturbance products to assess various forest ecology and resource management questions and issues.

IV. PROJECT ACTIVITIES AND OUTCOMES:

ACTIVITY 1: Process Landsat satellite imagery into a useable format

Description:

Although Landsat satellite data are now available free of charge from the USGS archive, processing is required to convert them into a useable format. We will acquire and process archived Landsat imagery that encompasses the Arrowhead region of Minnesota (~4.5 million acres). Specifically, this imagery will be obtained on a bi-monthly basis during all growing seasons over the last 40 years (1974-Present). The raw imagery will be processed to correct for atmospheric and geometric related errors to ensure image comparability. In total we will acquire and process approximately 500 images across the Arrowhead region. The total number of images acquired and processed will depend upon availability of suitable, cloud-free imagery. Our original plan was to process and analyze imagery for the entire forested region of Minnesota. However, due to the reduced level of funding recommended for this project we have decided to constrain the total area considered in this project. The reduction in acres processed is not directly proportional to the reduction in funding due to economies of scale. Specifically, the cost per acre of processing imagery decreases with an increase in total area. This is due to the fact that a baseline processing capacity (e.g., creating processing protocols and computer programming of processing algorithms) needs to be built regardless of the total area processed, and the cost associated with developing this baseline capacity is the same regardless of the total area of imagery processed.

Summary Budget Information for Activity 1:

ENRTF Budget: \$ 43,817
Amount Spent: \$ 43,817
Balance: \$ 0

Outcome	Completion Date
1. Complete Acquisition and Local Storage of Landsat Imagery	October 2015
2. Complete Pre-processing of Landsat Imagery	January 2016

Project Status as of February 2016:

After determining the appropriate time windows for growing season images using averaged NDVI curves (indicative of vegetation productivity) for each Landsat scene and assessing the quality of images for cloud cover and other initial sensor/data issues, we compiled lists of the large number of images within the required Landsat scenes needed to cover our study area and submitted the requested image lists to USGS. All images were downloaded, organized, and processed through an initial unpacking and quality assessment step. Although we attempted to choose images with low amounts of cloud cover, conditions in the Arrowhead region required us to include some images with clouds which had to be masked from the images during the atmospheric correction phase of pre-processing. Varying amounts of atmospheric correction were required depending on the Landsat sensor and the product available from USGS (data from more recent sensors are sometimes available as surface reflectance products processed by USGS, although this is not the case for earlier sensors which require additional pre-processing steps). Images from earlier sensors (the MSS series) also required additional geo-referencing efforts to ensure all images were properly aligned and resampled to match the spatial resolution of the TM and OLI images (30m x 30m pixels). We then calculated tasseled cap (brightness, greenness, and wetness) spectral indices for all images. As the Landsat sensors have some variation in the spectral bands and sensitivity through time, we calibrated the indices between sensors so that they are comparable through time and space, an important step in the pre-processing. Our final step will now be to create annual composites of the study area by averaging the inter-annual growing season indices for all scenes (thus producing annual stacks for the area representing the spectral trends through time to use in our next step of identifying disturbance/recovery trends using change algorithms, such as LandTrendr).

There has been a slight delay in the final composite step due to some additional work needed to complete the annual coverage of all areas. During initial processing, we discovered gaps in our yearly coverage of some areas during the early Landsat MSS sensors years (1972-1984). Adding these additional 12 years of historical information for our disturbance/recovery mapping efforts is extremely valuable for multiple applications. The problem is that some of the images we requested were only available from USGS in a less georeferenced form (L1G products), which makes them more difficult to directly include in the mostly automated steps appropriate for the other MSS images (L1T products). We have developed a solution for this issue using a semi-automated approach in which we first go through an initial geo-referencing step using manually detected tie points with a reference image in ArcGIS which does an initial correction on the image. The image is then ready to be entered back into the automated approach for final geo-referencing, tasseled cap calculations, calibrations, and ultimately, annual composites. The addition of this manual step has increased the expected processing time, but we expect to be finished with all pre-processing by the end of March.

Project Status as of September 2016:

After completing the pre-processing steps and exploring several available methods for compositing the image stacks into annual growing season spectral values, we created annual tasseled cap maps using the median of the input images per year. This compositing method further minimized undesirable noise and the appearance of seamlines between scenes within the annual maps. Activity 1 is now complete for the Arrowhead region for the years of 1972-2015.

Project Status as of February 2017:

Activity 1 is complete.

Project Status as of September 2017:

Activity 1 is complete.

Project Status as of February 2018:

Activity 1 is complete.

Final Report Summary:

The acquisition, pre-processing of all Landsat imagery, and ultimately the creation of harmonized annual composites of spectral indices, were the critical first products of our project which were integrated into all forest attribute modeling, disturbance mapping efforts, and assessments of forest trends. Our methodology for the harmonization of imagery from the different Landsat sensors, and the creation of our fitted annual composites, are published in our manuscript entitled “Extracting the full value of the Landsat archive: Inter-sensor harmonization for the mapping of Minnesota forest canopy cover (1973–2015)” (Vogeler et al. 2018). Our current goal is to secure future support to update these products beyond 2015 and develop semi-automated methods to facilitate annual additions to all mapping products in the future, leveraging recent developments in Google Earth Engine, a cloud computing platform. Our project is extremely unique in our inclusion of the more difficult to process MSS imagery, allowing for an additional decade of forest attribute and disturbance information to our time series analyses. This is significant for policy and management applications as we are now able to conduct more in-depth analyses of historic forest cover and its implications for future decision making. Any semi-automated methods developed through Google Earth Engine will be unable to replicate the inclusion of MSS data (which is still unavailable through that platform), but we hope to be able to add future years to our valuable dataset to ensure up to date forest maps so that the immense value of the products from this project can continue to stay current and relevant.

ACTIVITY 2: Disturbance database development and classification

Description:

We will employ a Landsat time series image analysis algorithm to detect forest disturbances from 1974-present. The basic approach involves using change detection to identify where disturbances occurred on the landscape as well as their severity and duration. In addition to detecting the disturbance, we will use semi-automated procedures to classify the type (i.e., cause) of each disturbance identified. At a minimum, disturbance type will be classified into the following categories (and subcategories): Harvesting (clear-cut, partial harvest, land use conversion) and Natural Disturbance (Insect Outbreak, Fire, and Blow Down). The semi-automated disturbance classification procedure attributes disturbance type to areas where disturbance has occurred based upon spatial and temporal patterns in the satellite image time series. An abrupt disturbance (e.g., clear-cut or high severity fire), for example, would likely be characterized by a sharp decline in the time series that is also spatially contiguous. Conversely, a more gradual or subtle disturbance (e.g., defoliating insect or low severity fire) will likely be characterized by a gradual decline in in the time series that is spatially variable (i.e., not spatially contiguous). Forest recovery rates, which are indicative of disturbance severity, will also be readily available for the time-series. Following disturbance identification and classification, historical information (e.g., FIA data, aerial photography, forest health surveys) and field data collected specifically for this project will be used to validate and assess the accuracy of the disturbance products.

Summary Budget Information for Activity 2:

ENRTF Budget: \$ 73,994
Amount Spent: \$ 73,994
Balance: \$ 0

Outcome	Completion Date
1. Disturbance patterns from 1974-2015 Identified	June 2016
2. Disturbances classified into primary categories	October 2016
3. Database with validated disturbance patterns over time completed	January 2017

Project Status as of February 2016:

No progress has been made on this Activity at this time due to the delay in image processing described above. We anticipate that outcome 1 will be completed on schedule or shortly thereafter, and all other outcomes will occur on schedule.

Project Status as of September 2016:

Although the initial calibration and other pre-processing steps of the project took longer than expected, we are close to being back on track for other scheduled outcomes. We have completed the first steps of the change algorithm, LandTrendr, which fits the annual composites with trend lines summarizing the spectral trajectories of each 30m x 30m image pixel. The process is called “segmentation” and the outputs are new annual spectral indices maps with the values smoothed to fit onto these trajectory lines. These new smoothed maps help to further reduce annual noise and serve as the input for the next tier of LandTrendr. The next step, referred to as “change labeling”, uses these fitted images to: 1) identify segments of disturbance/recovery and vertices between different segments; 2) group patches of pixels that may have experienced the same disturbance event; and 3) characterizes the disturbance/recovery events by their timing, magnitude of change (i.e. high severity vs. low severity), and duration (i.e. abrupt vs. slow). Characteristics of the disturbance and recovery events within patches aid in the classification of disturbance agent (i.e. harvest vs. insect infestation). We have begun to test the change labeling LandTrendr algorithm on a subset of the Arrowhead region, although there is much calibration of parameters still needed before we are able to apply the algorithm to the whole region.

In addition to serving as inputs to the change labeling portion of LandTrendr, we also incorporated the smoothed annual spectral images from the segmentation procedure into historical canopy cover mapping efforts. We estimated canopy cover at a subset of FIA plots locations using 4-band NAIP imagery, then extracted tasseled cap predictors from the corresponding year of segmentation images to create statistical models of forest cover. We are currently working on selecting the best model and preparing the other years of imagery, for which we will then apply the model to create maps of annual estimated canopy cover from 1972-2015. These historical canopy cover maps may help identify conversion of land to/from forest cover types through time, estimate trends of canopy cover change due to disturbance/recovery, and characterize an important habitat component for wildlife habitat monitoring efforts through time.

Project Status as of February 2017:

After creating an extensive training canopy cover dataset from NAIP imagery across all land cover types, testing multiple statistical modeling approaches and Landsat time series along with static predictors to find the best model for minimizing prediction error, we have selected our final models for delineating forest vs non-forest and for predicting continuous canopy cover across the entire Arrowhead region. We are currently applying these models to the stacks of smoothed/fitted tasseled cap outputs from LandTrendr time series stacks to create annual forest mask and canopy cover models across all land cover types from 1972-2015. Predictive mapping across large spatial areas as well as across 44 years of imagery is computer intensive and thus takes a significant amount of time, although these products should be complete by the end of February. Immediate applications of the products are already being outlined in addition to their use as valuable inputs for the identification and interpretation of change and recovery within our forest disturbance mapping and classifications.

As we finish these forest attribute maps which will also aid in the disturbance mapping and classification, we continue to test the LandTrendr change algorithm across the Arrowhead region. There are multiple parameters within LandTrendr which must be calibrated for specific study areas as well as setting thresholds within the outputs for designating patches as true change. While we are behind in the timeline for Activity 2, we feel strongly that these additional products and calibration steps will be vital in creating quality change detection

products as well as providing important canopy cover information for wildlife habitat modeling, forest stand characterization, and monitoring trends in overall forest area across the last 44 years. Incorporating the early Landsat sensors into time series stacks to add an additional 12 years of forest information, such as we are doing here, is rarely done and thus timelines for spectral and spatial calibrations throughout the process were difficult to predict.

We have begun to turn our attention to creating our forest disturbance agent database for training and validation of change agent classifications. We have developed methods for delineating harvest boundaries within Google Earth historic imagery, we have compiled the FIA plot data for our study region and are exploring attributes which may be extracted from this extensive data base, and we continue to explore new avenues for acquiring accurate spatial information for forest disturbance agents and recovery characteristics.

Project Status as of *September 2017*:

We have completed the annual forest mask and canopy cover maps including analyzing forest cover trends across the >40 year study period, and the associated manuscript has been prepared and submitted to Remote Sensing of Environment. Initial reviews were favorable and we are currently addressing manuscript revisions. We are now exploring the change detection products from initial runs of the second step of the LandTrendr algorithm. We have focused our efforts on the “fast” disturbance patches to begin with, which are disturbances that have durations of 4 years or less (i.e. harvest, fire, and wind blow). Change patch predictors have been extracted for these initial patches including LandTrendr derived pre- and post- disturbance spectral values, magnitudes of change, along with additional spatial predictors. We are working to compile the “fast” agent training data base and have begun to explore classification modeling using preliminary training data, although we continue to explore additional predictors and training data sources. We have identified preliminary versions of “slow” disturbance patch maps (i.e. those resulting from insects and other agents of forest decline), although change parameters for defining the final patches are still being calibrated for the Arrowhead region’s disturbance patterns and agents. We are primarily utilizing historic photos in google earth and the annual forest health aerial surveys for the extraction of fast and slow agent classification training data. In August, we conducted field visits to various forest and disturbance types in the study region as well as meeting with foresters from the Chippewa National Forest, Superior National Forest, and UMN Cloquet Forestry Center to discuss local disturbance patterns, future applications of mapping products, and the needs of the on-the-ground forest managers.

Project Status as of *February 2018*:

While initial change detection products and models classifying the agents of change for fast disturbances (i.e. harvest, fire, wind, and land conversions) have been created, we continue to work to improve the quality of these products through a multi-iterative approach common for Landsat time series studies. In this multi-iterative approach, initial change patches are identified, agent training data is collected, predictive metrics from the time series spectral indices as well as other ancillary data are created, classification models are selected, and classified change maps are produced. Within Landsat time series analysis methodologies, after such a “run” is finished, the products are thoroughly assessed for quality and accuracy, potential areas for improvement are identified, and steps are taken to either add needed training data or expand predictors to then proceed with a new “run” of the data and mapping process. These steps are repeated until a level of quality is achieved that meets projects goals, at which time finalized change maps undergo independent validation efforts which are presented with the mapping products and within publications so that the strengths and limitations of the data are well understood by all future users. We have currently undergone several of these “runs” of our classified change maps, and we are working towards finalizing our models and maps of fast disturbances to proceed with validation efforts. Coding and workflows have also been created for analyzing the trends from the classified change maps so that when finalized products are created, the framework is in place to conduct further trends analyses to include in planned publications.

Final Report Summary:

Among the foundational mapping products created through this project were 43 years of annual forest mask and percent canopy cover maps for the entire state of Minnesota. The canopy cover products have been integrated into several application projects as well as serving as the basis for assessing forest trends in Minnesota across 4 decades within the publication in Remote Sensing of Environment (Vogeler et al. 2018). The canopy cover maps have also played an important role as model inputs for classifying the agent of change in our disturbance mapping efforts.

We have now also completed the mapping of the most recent fast disturbances dating back to 1974 across the Laurentian Mixed Forest Province, classified by the agent of change, also an unprecedented temporal and spatial data set for Minnesota. We expanded beyond the Arrowhead region to better match the focal region of many of the proposed application collaborations. After exploring several mapping approaches, we decided to focus on fast disturbance events (durations of 4 years or less) in our change mapping methods. We believe that these abrupt change events include the majority of the natural and anthropogenic disturbance agents in this region, and are able to be identified through aerial photos and existing spatial data for the creation of training and validation data sets. We used a minimum mapping size of 11 pixels, as the focus of this project was on events that impact at least a large portion of a forest stand. This size limit also increased our confidence in our change detection by excluding smaller events that may have been captured in error as a product of spectral noise. Out of the spectral indices created in this project, tasseled cap wetness (TCW) has been identified as the most indicative of forest canopy structure by previous studies. Therefore, we utilized thresholds of TCW change representative of moderate- to high- intensity changes to identify our disturbance patches. We converted initial raster based change information to polygons for the next steps in map creation and disturbance agent classifications.

Agents included in the classification represent the common moderate- to high-intensity natural and anthropogenic disturbance agents characteristic of the study region, including: harvest, land conversion, fire, wind/weather, flooding, and an “other natural” category for rare natural events not represented by the other classes. Training data of known disturbance events for a sample of the change polygons was opportunistically compiled using a combination of google earth historic imagery, the national Monitoring Trends in Burn Severity fire perimeter maps, online historical records of large natural disturbance events often reported by county which could be used in conjunction with photos for identifying change areas, and personal communications with local land managers for more recent large-scale natural events. Using a sample of known events as our training data set, we employed a random forest classification approach to predict the agent for all disturbance polygons across the study area. We mapped these classified agents to create a map of the most recent fast disturbances across the study area dating back to 1974.

To validate the accuracies of the initial change/no-change maps, we created a random sample of points within forests across the study area with buffers of 3x3 pixels. We assessed if changes had occurred within the validation buffers, and whether those events were captured by our change detection methodologies. This also allowed us to evaluate potential limitations and/or biases in our maps to later relay to map users. Within the validation efforts, we utilized historical imagery within google earth and the Minnesota Historical Aerial Photos Online database (<https://www.lib.umn.edu/apps/mhapo/>), as well as any additional spatial information we had compiled for the disturbance agent modeling work. We choose to focus on stand replacing events that we could validate with some confidence and which were the major focus of our project. We therefore excluded flooding (which was often the result of change in moisture in overlapping forest and riparian non-forest areas and did not always result in stand mortality), and the “other natural” polygons (which were associated with more rare within stand natural events such as tent caterpillar outbreaks, which did not always result in large changes in canopy and which were difficult to confidently validate in older black and white historic photos). While we were able to identify polygons for training purposes within the flooding and “other natural” categories, we did so

opportunistically for areas with known events. Identifying these events with confidence across larger areas and back in time within black and white historical photos proved to be more difficult. Our validation efforts thus focused on a reduced map of change polygons classified as harvest, land conversion, fire, or wind/weather. Within the reduced map, if we were not able to confidently assign a change or no-change classification (for instance, historical photos were not available for a given point), we excluded this validation plot from our data.

Our final maps of the most recent fast disturbances had an overall change/no-change classification accuracy of 92%, with a TPR (sensitivity) of 0.859, and a TNR (specificity) of 0.960 for moderate- to high-intensity stand replacing events. Our final random forest classification model for the agent of change also had an overall classification accuracy of 92% (as calculated from out-of-bag error rates), with class accuracies ranging from 78% to 96%, where the harvest class had the highest accuracies and fire had the lowest classification success.

ACTIVITY 3: Development of spatial descriptors and application of findings

Description:

Statistical descriptors of the disturbance patterns will be derived from the disturbance database. These statistical descriptors will be integrated into models assessing disturbance impacts on the current status of two critical resources: wildlife habitat (specifically moose) and wood resources (specifically the amount, type, and quality). In these example applications, we will compare existing moose population demography data and forest inventory data with disturbance descriptors in a modeling framework. This modeling framework will provide a quantitative assessment of how the temporal and spatial configurations of specific disturbance types can either enhance or degrade the sustainability of the resources (Figure 1). Through this process we will identify management responses that will sustain and or improve forest resources under future disturbances. Results quantifying the impacts of disturbance dynamics on the health and resilience of forest resources will be summarized in public project reports and conveyed to forest managers through outreach activities.

Summary Budget Information for Activity 3:

ENRTF Budget: \$ 82,189
Amount Spent: \$ 82,189
Balance: \$ 0

Outcome	Completion Date
1. Develop statistical descriptors of disturbance patterns	August 2017
2. Model impacts of disturbance dynamics on forest resources	January 2018
3. Publish project summaries and conduct outreach activities	June 2018

Project Status as of February 2016:

Work on this activity will commence after completion of Activity 2.

Project Status as of September 2016:

Work on this activity will commence after completion of Activity 2.

Project Status as of February 2017:

Work on this activity will commence after completion of Activity 2.

Project Status as of September 2017:

Work on this activity will commence after completion of Activity 2.

Project Status as of February 2018:

Our first manuscript from this project, titled “Extracting the full value of the Landsat archive: Inter-sensor harmonization for the mapping of Minnesota forest canopy cover (1973-2015)”, has been accepted for publication within Remote Sensing of Environment. This manuscript presents many of the initial methods of our project for the harmonization of the Landsat time series stacks, creation of spectral indices, and the modeling and mapping of annual statewide canopy cover and trends through time. As we work to finalize our change detection products classified by agent and associated validation efforts, we are outlining additional forest trend analyses and associated manuscripts to publish findings of this project.

In January 2018, we presented our project methods, forest attribute products, and initial forest disturbance trend results at several MN meetings, including an invited webinar to members of MN DNR, forest service, and UMN affiliates, and presenting at the 2018 SFEC Forestry and Wildlife Research Review meeting at the Cloquet Forestry Center. We also conducted smaller meetings with members of MN DNR, forest service, and UMN, to discuss potential collaborations and applications of our mapping products and project results. All of these meetings displayed a wide breadth of interest from a variety of research and management agencies in the use of our forest attribute and change detection products for a range of forest, wildlife, water quality, and monitoring applications. Several collaborations are also now underway for the use of these products in the assessment of wildlife habitat and the impacts of management and other drivers of landscape changes through time.

Final Report Summary:

Through our annual mapping of forest/non-forest masks and percent canopy cover, we were able to evaluate forest trends across the state of Minnesota dating back to 1973. These trends were presented in our Remote Sensing of Environment publication (Vogeler et al. 2018). Generally, we observed a significant, although slight, positive trend in forest area across the state throughout the time period of the study (1973-2015). It should be noted that this trend does not account for changes in forest types or stages of structural development, and may include some areas not commonly classified as forests (e.g., shrub-scrub wetlands). Trends were also evaluated within specific Minnesota ecological provinces, where the significant positive trend in forest area was maintained across all provinces. We also assessed temporal trends of forest area within five canopy cover classes estimated using our percent canopy cover maps. Across the 43 years of the study, all provinces experienced significant growth in the land area within the two highest cover classes (50-74% and 75-100%), with the exception of a non-significant trend for the 50-74% class within the Tallgrass Aspen Parkland. The greatest growth in the two highest cover classes occurred within the Laurentian Mixed Forest province.

The base spectral composites utilized within all mapping products within this project, as well as the forest attribute and disturbance mapping products themselves, have been (or are currently being) incorporated into several wildlife, forest resources, and water quality assessment application projects, although we continue to establish additional collaborations and explore new applications for the data. Some notable examples include:

- Early in the project, we utilized our fitted Landsat-derived spectral indices to create bi-annual land cover maps for the Arrowhead region (1999-2015) as requested by University of Minnesota moose biologist, James Forester, to facilitate the assessment of moose habitat use and movements as impacted by land cover types and changes through time. This data is being used to develop predictive models of moose population dynamics to address ongoing questions related to long-term moose viability in Minnesota.
- We are also collaborating with wildlife researchers at the University of Minnesota-Duluth and the Natural Resources Research Institute to assess the impacts of harvest characteristics (e.g. quantity and spatial arrangement of retained canopy during harvest activities) on avian and small mammal communities through the incorporation of our canopy cover products and disturbance map. The study includes an array of aspen dominated stands across the Arrowhead region representing a chronosequence of harvest years. We are utilizing our disturbance products to assign the initial year of harvest

activity, as well as assessing change intensities. Our annual canopy cover products are facilitating the characterization of stands prior to harvest, changes in canopy after timber removal, and the recovery of canopy at the time of wildlife surveys. Ultimately, this data is being used to determine the suitability of Minnesota's forest management guidelines at maintaining wildlife populations following forest harvesting.

- Recently, we provided initial harvest disturbance maps to the Minnesota consulting firm, RESPEC, for incorporation into a tool for assessing best management practices and impacts on water quality. Below is a write-up pertaining to the utilization of our disturbance mapping products provided by our RESPEC liaison, Paul Marston:

“RESPEC Consulting was hired by the Minnesota Pollution Control Agency (MPCA) to add forestry best management practices (BMPs) to the existing Scenario Application Manager (SAM) user interface for the Hydrologic Simulation Program-Fortran (HSPF) models. The MPCA has invested in HSPF to model the entire state of Minnesota at the HUC-8 watershed level. SAM was developed to allow watershed professionals a tool to access the data from the HSPF models as well as run scenarios using the model information. SAM allows users to apply BMPs to certain land use classes and rerun the model to determine the water quality implications. To incorporate forestry BMPs into SAM, a key data need was quantifying the area of harvest on a year to year basis. The harvest area data was the foundation for determining what areas within the model framework could have forestry BMPs applied. The disturbance product provided by Dr. Falkowski and his team was critical as it was the only data available that identified specific agents of forest canopy disturbance. Using Dr. Falkowski et al.'s work gave the project team strong confidence that a critical data requirement for our methodology was met. Other forest disturbance data available grouped all agents of forest canopy change together, providing a dataset with areas not specific to harvesting areas. This data was used in areas of the state that Dr. Falkowski et al.'s data did not cover. Our hope in the future, would be to have statewide versions of Dr. Falkowski et al.'s disturbance products to quantify the harvest area for the entire state of Minnesota, which would ensure we are using the most accurate data available for a critical component of our project.”

Paul Marston
Watershed Scientist
RESPEC
Paul.Marston@respec.com
651-305-2278

Among the proposed outcomes of Activity 3, was the publishing of project summaries and conducting outreach activities. In addition to the already published manuscript entitled “Extracting the full value of the Landsat archive: Inter-sensor harmonization for the mapping of Minnesota forest canopy cover (1973–2015)” (Vogeler et al. 2018), we are currently working towards the submission of a manuscript summarizing our most recent fast disturbance mapping work and associated applications. We hope to continue to publish work related to the data products from this project as existing collaborations pertaining to the applications of the mapping products move forward.

Project team members have also been active in the presentation of project methods and preliminary results to a variety of audiences and potential users of the data over the last few years. These included two invited webinars, one to the MN Department of Natural Resources and UMN affiliates, the other to the USDA Forest Service's Forest Inventory and Analysis (FIA) Research and Techniques Band (recording available at: <https://usfs.adobeconnect.com/prjhzov1f5fi/>), as well as a department seminar within the Department of

Ecosystem Science and Sustainability at Colorado State University in Fort Collins, CO. We also participated in the 2018 Annual Forestry and Wildlife Research Review at the Cloquet Forestry Center, with our presentation titled "A foundational data set characterizing historic forest attributes and disturbance patterns". Our work has even reached the international forest remote sensing community through our presentation of early project methodologies at the bi-annual ForestSAT conference in Santiago, Chile in 2016 (attended using non-ENRTF funds). We will once again present to this international community at this year's ForestSAT held in College Park, MD in October 2018. In addition to the many official presentations, our team has also met with countless researchers and managers around Minnesota, representative of a number of state and federal agencies as well as UMN affiliates, including site visits with local managers at the Chippewa National Forest, Superior National Forest, and the Cloquet Forestry Center.

PROJECT PUBLICATIONS:

Vogeler, J.C., J.D. Braaten, R.A. Slesak, M.J. Falkowski. 2018. Extracting the full value of the Landsat archive: Inter-sensor harmonization for the mapping of Minnesota forest canopy cover (1973 -2015). *Remote Sensing of Environment*, 209, 363-374.

PROJECT PRESENTATIONS:

Vogeler, J. C., R. A. Slesak, and M. J. Falkowski. 2018. Characterizing forest change dynamics in Minnesota for forest and wildlife science applications (1975-2015). USDA Forest Service Forest Inventory and Analysis Research Techniques Band, Invited Webinar, 26 April 2018. Recording available at: <https://usfs.adobeconnect.com/prjhzov1f5fi/>

Vogeler, J. C., R. A. Slesak, and M. J. Falkowski. 2018. Characterizing Forty Years of Forest Change: Applications in Forest and Wildlife Science. Department of Ecosystem Science and Sustainability Seminar, Colorado State University, Fort Collins, CO, 18 April 2018.

Vogeler, J. C., J. Braaten, R. A. Slesak, and M. J. Falkowski. 2018. A foundational data set for >40 years of forest characterization and change detection. Minnesota Department of Natural Resources Webinar, St. Paul, MN, 9 January 2018.

Vogeler, J. C., J. Braaten, R. A. Slesak, and M. J. Falkowski. 2018. A foundational data set characterizing historic forest attributes and disturbance patterns. *Forestry and Wildlife Research Review*, Cloquet, MN, 11 January 2018.

Vogeler, J. C., J. Braaten, R. A. Slesak, and M. J. Falkowski. 2016. Mapping historical canopy cover change and recovery using Landsat time series imagery (1972-2015). *ForestSAT*, Santiago, Chile, 16 November 2016.

V. DISSEMINATION:

Description:

The final product of this project will be a digital geospatial database characterizing disturbance trends and types across the forested regions of Minnesota. In addition we will prepare an interpretative report detailing case studies that demonstrate how the geospatial disturbance database can be used to assess the impacts of historic disturbances on the resilience of three key forest resources: water quality, wildlife habitat, and wood fiber. We will work in conjunction with the Minnesota Department of Natural Resources to make the geospatial database publically available for download from the GIS Data Deli website (<http://deli.dnr.state.mn.us>). The interpretative report will be made available on the Internet as a Department of Forest Resources Staff Paper Report. In addition, several manuscripts will be written based on this research and submitted for publication in peer-reviewed journals. A fact sheet summarizing principal findings of this project will be distributed to LCCMR members and legislators at the state and federal level. Results will be presented at state and national forest

management and forest health conferences, and notably to agency and individual participants in the Sustainable Forests Education Cooperative. All reports and publications from this project will be made available via the Department of Forest Resources web site (www.forestry.umn.edu).

Project Status as of February 2016:

Work on this activity will commence after completion of Activity 3

Project Status as of September 2016:

Work on this activity will commence after completion of Activity 3.

Project Status as of February 2017:

Work on this activity will commence after completion of Activity 3.

Project Status as of September 2017:

Work on this activity will commence after completion of Activity 3.

Project Status as of February 2018:

Work on this activity will commence after completion of Activity 3 (although see above section for current publication work and recent presentations/meetings).

Final Report Summary:

As we finish packaging our mapping products, we are compiling the necessary metadata for posting our annual canopy cover and most recent fast disturbance maps to the MN DNR GIS Data Deli website, which we anticipate will be completed in the next few months. We will also work to post all publications and project reports to UMN Department of Forest Resources web site as they become available. A fact sheet summarizing principal findings of this project is included in the supplementary materials, along with several visuals related to our work and results, and our publication. In addition to the publication and presentations listed above, we are working on an additional manuscript summarizing the most recent fast disturbance mapping work and associated applications and continue to present our project results and mapping products to various research and management audiences and other potential future users of the data. In addition, project partner Dr. Robert Slesak is actively exploring additional applications of this dataset through his work at the MN Forest Resources Council and related activities.

VI. PROJECT BUDGET SUMMARY:

A. ENRTF Budget Overview:

Budget Category	\$ Amount	Overview Explanation
Personnel:	\$ 193,500	-Salary and fringe (0.336) for three years for Falkowski - PI; 0.083 FTE each year (0.25 FTE over entire project) -Salary and fringe (0.336) for a post-doctoral researcher; 1.0 FTE for 2.5 project years (2.5 FTE over entire project) -Salary and fringe (0.336) for a technician; 0.083 FTE for 2 years (0.16 FTE over entire project).

Equipment/Tools/Supplies:	\$ 1,500	Equipment and supplies include \$1,500 for data hard drives for storing the satellite data and archiving the final geospatial disturbance database.
Travel Expenses in MN:	\$ 5,000	This money will be used to pay for mileage (\$3,750) and lodging (\$1,250) for researchers when performing validation of the disturbance dataset.
TOTAL ENRTF BUDGET:	\$ 200,000	

Explanation of Use of Classified Staff: N/A

Explanation of Capital Expenditures Greater Than \$5,000: N/A

Number of Full-time Equivalents (FTE) Directly Funded with this ENRTF Appropriation: 2.91

Number of Full-time Equivalents (FTE) Estimated to Be Funded through Contracts with this ENRTF Appropriation: N/A

B. Other Funds: N/A

VII. PROJECT STRATEGY:

A. Project Partners:

In addition to the project leader, Michael Falkowski other project partners are included below.

Dr. Alan Ek, Department of Forest Resources – University of Minnesota (not receiving funding). Role: Dr. Ek will serve as a liaison to several county forestry departments in the State of Minnesota who will be able to assist with database validation by providing disturbance datasets, and will ultimately be end users of the final disturbance database.

Dr. Joe Knight, Department of Forest Resources – University of Minnesota (not receiving funding). Role: Dr. Knight will assist with some remote sensing aspects of this project primarily by providing feedback to the post-doctoral researcher.

Dr. Matthew Russell, Department of Forest Resources & and Extension– University of Minnesota (not receiving funding). Role: Dr. Russell will assist with characterizing insect and disease related disturbances in the final database. In addition he will assist in project dissemination via his role in the University Extension program.

Dr. Linda Nagel, Department of Forest Resources – University of Minnesota (not receiving funding). Role: Dr. Nagel will assist with characterizing harvest related disturbances in the final database

Dr. Robert Slesak, Minnesota Forest Resources Council. Role: Dr. Slesak will assist with the water quality related applications of the disturbance database. Dr. Slesak will also serve as a liaison to Forest Resource Council members who will ultimately be end users of the final disturbance database. In addition he will integrate efforts of this project with other projects he’s currently working on in conjunction with the Minnesota Department of Natural Resources Resource Assessment and Wildlife groups.

Collaborators will include the Minnesota Department of Natural Resources Resource Wildlife and Assessment groups, the Superior National Forest, University of Minnesota Extension, and several counties in northern Minnesota.

B. Project Impact and Long-term Strategy:

Due to the multiple disturbance threats (e.g., insect outbreaks, fire, conversion to agriculture, and climate related stress) facing Minnesota's forest resources, as well as the fact these threats will only increase under projected climate change, there is a critical need for datasets that can be used to assess the impacts of disturbance on the long term sustainability of Minnesota's forest resources. In addition, understanding how past disturbances have influenced current forest resources is essential to improving and sustaining future resource conditions under existing and eminent threats. This 3 year project will develop a foundational dataset characterizing historic forest disturbance dynamics -and related resource impacts- which will be a powerful tool for identifying threshold disturbance patterns that impact multiple forest resources across Minnesota. This foundational dataset will allow us to evaluate how disturbance and landuse configuration over the past 40 years have influenced the current status of forest resources and help to identify management responses that improve and sustain forest resources into the future, and will ultimately guide forest management response aimed at avoiding or mitigating persistent detrimental impacts of forest disturbance on forest resources. For example, forest managers will be better equipped to strategically plan disturbance mitigation practices where risks to forest resources are high, or manipulate disturbed areas to enhance forest response in a manner beneficial to multiple resources. Given the long-term nature of forest disturbance dynamics and associated management, we will link our work with on-going work in the MNDNR resource assessment office that is focused on using similar technology to understand future disturbance impacts. We also plan to explore additional funding opportunities from federal sources such as NASA, the National Science Foundation, and the US Forest Service to build upon and extend this work into the future.

C. Funding History: N/A

VIII. FEE TITLE ACQUISITION/CONSERVATION EASEMENT/RESTORATION REQUIREMENTS: N/A

IX. VISUAL COMPONENT or MAP(S):

See attached Figure 1

X. RESEARCH ADDENDUM: N/A


XI. REPORTING REQUIREMENTS:

Periodic work plan status update reports will be submitted no later than September 2015, February 2016, September 2016, February 2017, September 2017, February 2018. A final report and associated products will be submitted between June 30 and August 15, 2018.


Supplementary Material 1. Project Summary Fact Sheet

- We were able to combine data from all Landsat sensors, allowing for the mapping of >40 years of forest attributes and forest change detection in Minnesota.
- Use of smoothed time series composites highlighted the value of using a time series context (as opposed to 2 date methodologies) and a segmentation fitting approach for more stabilized land cover and forest attribute mapping through time, as well as improved change detection.
- Our annual forest mask and canopy cover mapping products provided the opportunity to observe forest area and cover trends across the state from 1973-2015. We observed a significant, although slight, positive trend in forest area across the state throughout the time period of the study, although we should note that this trend does not account for changes in forest types or stages of structural development, and may include some areas not commonly classified as forests (e.g., shrub-scrub wetlands).
- All ecological provinces experienced significant growth in land area within the two highest cover classes (50-74% and 75-100%), with the exception of a non-significant trend for the 50-74% class within the Tallgrass Aspen Parkland. The greatest growth in the two highest cover classes occurred within the Laurentian Mixed Forest province.
- Our change detection approach captured moderate- to high-intensity stand events with overall accuracy of 92% for differentiating areas of change vs. no-change across Laurentian Mixed Forest Province.
- We were able to further classify the most recent fast disturbance patches into 6 change classes representative of the fast anthropogenic and natural disturbance agents common to the study area (e.g., forest harvest, fire, wind disturbance, etc.), with class accuracies ranging from 78-96% with an overall accuracy of 92%.
- Forest attribute and disturbance products from this project provide unprecedented spatial and temporal information for a variety of Minnesota forest resources applications.
- We utilized our fitted Landsat-derived spectral indices to create bi-annual land cover maps for the Arrowhead region (1999-2015) as requested by University of Minnesota moose biologists to facilitate the assessment of moose habitat use and movements as impacted by land cover types and changes through time. This data is being used to develop predictive models of moose population dynamics to address ongoing questions related to long-term moose viability in Minnesota.
- We are currently collaborating with wildlife researchers at the University of Minnesota-Duluth and Natural Resources Research Institute to assess the impacts of harvest characteristics (e.g. quantity and spatial arrangement of retained canopy during harvest activities) on avian and small mammal communities through the incorporation of our canopy cover products and disturbance map.
- We have provided harvest disturbance maps to the Minnesota consulting firm RESPEC, under contract to the MN PCA, for incorporation into a watershed planning tool for assessing best management practices and impacts on water quality.


Landsat Time Series Analyses for Forest Research and Management




Two Date Change Detection
VS
Time Series Trends




By looking at a longer time series of images, we can gain more insight into forest trends as well as creating more stabilized products by having the context of the time series.



Supplementary Material 3. Project mapping product extents. Annual maps of canopy cover were created at the Minnesota state extent (1973-2015), while disturbance mapping and change agent classification efforts were focused on the Laurentian Mixed Forest Province (with an additional 15km buffer).




Supplementary Material 4. Most Recent Fast Disturbance Map for the Laurentian Mixed Forest Province with 3 example inset areas. Inset A is part of the Chippewa National Forest, inset B represents a mixed ownership landscape, and inset 3 falls within the Boundary Waters Canoe Wilderness Area.



Supplementary Material 5. Visual example of a wildlife application of our data products where we utilized our disturbance map to identify the initial year of harvest activities for study stands and characterized the amount and spatial arrangement of retained canopy and vegetation regrowth following harvest from our canopy cover data sets. This project is a collaboration with wildlife researchers from UMN-Duluth and Natural Resources Research Institute.

Leave Tree Project Objective: Evaluate the impact of the amount and distribution of retained canopy during harvest activities on wildlife communities, as well as quantify canopy recovery since harvest at time of wildlife surveys.


2004 Harvest




2007 Harvest




2014 Harvest




Immediate Post-Harvest



2015 Vegetation Regrowth



Canopy Cover (%)



Supplementary Material 6. Publication entitled “Extracting the full value of the Landsat archive: Inter-sensor harmonization for the mapping of Minnesota forest canopy cover (1973–2015)”. Published in Remote Sensing of Environment (Vogeler et al. 2018). Delivered as pdf with this report.



**Environment and Natural Resources Trust Fund
M.L. 2015 Project Budget**

Project Title: Foundational Dataset Characterizing Historic Forest Disturbance Impacts
Legal Citation: M.L. 2015, Chp. 76, Sec. 2, Subd. 03q
Project Manager: Michael J. Falkowski
Organization: University of Minnesota
M.L. 2015 ENRTF Appropriation: \$ 200,000
Project Length and Completion Date: 3 Years, June 30, 2018
Date of Report: 08/17/2018

ENVIRONMENT AND NATURAL RESOURCES TRUST FUND BUDGET	Activity 1 Budget	Amount Spent	Activity 1 Balance	Activity 2 Budget	Amount Spent	Activity 2 Balance	Activity 3 Budget	Amount Spent	Activity 3 Balance	TOTAL BUDGET	TOTAL BALANCE
BUDGET ITEM	Process Landsat satellite imagery into a			Disturbance database development and			Development of spatial descriptors and				
Personnel (Wages and Benefits)	\$42,317		\$42,317	\$71,494	\$0	\$70,494	\$79,689	\$0	\$79,689	\$193,500	\$0
Michael Falkowski, Project Manager: \$8,684 (66.4% Salary, 33.6% benefits); 0.083 FTE each year (0.25 FTE over entire project).		\$1,430			\$2,576			\$24,746			
Jody Vodeler, Postdoctoral Researcher: \$167,000 (66.4% Salary, 33.6% benefits); 1.0 FTE each year for 2.5 years (2.5 FTE over entire project).		\$41,403			\$69,494			\$49,120			
Reinhardt, Jason, Technician: \$6,948 (66.4% Salary, 33.6% benefits); 0.083 FTE for 2 years (0.16 FTE over entire project)								\$6,702			
Equipment/Tools/Supplies											
Equipment and supplies include \$1,500 for data hard drives for storing the satellite data and archiving the final geospatial disturbance database.	\$1,500	\$984								\$1,500	\$0
Travel expenses in Minnesota											
This money will be used to pay for mileage (3,750) and lodging (1,250) for researchers when performing validation of the disturbance dataset.				\$2,500	\$1,924	\$2,500	\$2,500	\$1,621	\$2,500	\$5,000	\$0
COLUMN TOTAL	\$43,817	\$43,817	\$0	\$73,994	\$73,994	\$0	\$82,189	\$82,189	\$0	\$200,000	\$0